

# HUMAN EXPOSURES TO INORGANIC MERCURY

Among the major sources of human exposures to inorganic mercury are dental amalgams, mercury spills, and occupational exposures. In addition to dental amalgams, inorganic mercury has a variety of other medical and dental uses. Inorganic mercury is also widely used in industrial processes and consumer products. The physical properties of mercury make it an ideal material for use in measuring devices, switches, fluorescent lamps, and in lab and pharmaceutical chemicals.

For detailed information on exposures to and the health effects of inorganic mercury, see EPA's *Mercury Study Report to Congress*<sup>1</sup> and the Agency for Toxic Substances and Disease Registry's *Toxicology Profile on Mercury*.<sup>2</sup>

## MEDICAL AND DENTAL USES OF INORGANIC MERCURY

**Medical uses.** Hospitals and the health care industry are large users of mercury. Medical waste incinerators are the fourth largest known source of mercury emissions into the environment in the United States.<sup>1</sup>

Some of the most common uses of mercury in hospitals include: blood pressure monitors, thermometers, esophageal dilators, feeding tubes, and laboratory chemicals.

**Dental offices.** Mercury is a primary ingredient in mercury-silver amalgam dental fillings. Inorganic mercury contamination of dental offices can occur unless a great deal of care is taken in preparation of these amalgams. Once in place in the tooth, these amalgams may release mercury. Following chewing of food, gum chewing,<sup>3</sup> or "grinding" of teeth,<sup>4</sup> inorganic mercury (as mercury vapor) is released from the filling and can be measured in the mouth with special instruments. Some of this mercury is swallowed and accumulates in tissues.<sup>5-8</sup> A correlation has been found between inorganic mercury in human breast milk and mercury-silver dental amalgams in the mother.<sup>9-11</sup> There is an ongoing controversy over whether people experience adverse effects from the amount of mercury released by mercury-silver amalgam restorative materials.<sup>12-14</sup>

**Folk healing practices.** Mercury is used by some groups in the US for healing purposes. Medicinal uses include swallowing mercury and sprinkling it in bath water. (Some groups use mercury in their religious practices; spiritual

uses include wearing amulets containing mercury or sprinkling it around a room or bed for good luck.)

## MERCURY IN CONSUMER PRODUCTS

Mercury is found in numerous consumer products. In addition to fever thermometers and the mercury switches that regulate our thermostats, mercury is used in many other common products, including measuring devices, fluorescent lamps, and pilot light sensors. (Mercury-containing sensors are found in some gas-fired appliances that have pilot lights, such as stoves, water heaters, and furnaces.) Mercury is also found in certain laboratory and pharmaceutical chemicals such as reagents.

**Measuring devices.** Many thermometers and barometers contain elemental mercury. Most mercury fever thermometers are extremely fragile and can be easily broken, allowing mercury to escape. Accidental mercury releases in the home pose a significant poisoning risk for children.

**Switches.** Mercury switches are used in a number of applications. Many home thermostats use mercury switches to control heating systems. Silent light switches use a mercury switch rather than a mechanical one to turn on and off lights. In automobiles, mercury is found in switches that turn on trunk and hood lights and in some anti-lock breaking systems.

**Lighting.** Fluorescent lamps also contain small amounts of mercury. The industry has reduced the amount of mercury in lamps since the mid 1990s; although the amount in each lamp is relatively low, some of the mercury is immediately released to the atmosphere once a lamp is broken, so it is critical that used lamps are managed appropriately.

## OCCUPATIONAL EXPOSURE TO INORGANIC MERCURY

In the 19th century, hat makers' exposure to mercury used as a fixative may have given rise to the familiar term "mad as a hatter." Occupational exposure to mercury may occur in hospitals, laboratories, dental practices, and in industrial workplaces, particularly in the chemical and drug manufacturing industries.<sup>15</sup> Workers may be exposed to mercury during the manufacture of:

measuring instruments, mercury arc lamps, mercury switches, fluorescent lamps, mercury boilers, mirrors, electric rectifiers, electrolysis cathodes, pulp and paper, zinc carbon and mercury cell batteries, dental amalgams, antifouling paints, explosives, and disinfectants. Workers in the fur processing industry are also at risk for exposure to inorganic mercury. Other industrial uses of mercury include metallic mercury, mercury salts, mercury catalysts (used in making urethane and epoxy resins), mercury fulminate, Millon's reagent, chlorine and caustic soda, pharmaceuticals, and antimicrobial agents.<sup>16</sup>

**Jeri Weiss, MS**  
**Luke Trip**  
**Kathryn R. Mahaffey, PhD**

Ms. Weiss is an Environmental Protection Specialist with Region I, US Environmental Protection Agency (EPA), Boston. Mr. Trip is the Manager, Heavy Metals Issues, Environment Canada, Quebec. Dr. Mahaffey is a Senior Scientist with the National Center for Environmental Assessment, EPA, Washington, DC.

**FOR MORE INFORMATION:**


Jeri Weiss, EPA, 1 Congress Street (CME), Suite 1100, Boston MA 02114; tel. 617-918-1568; fax 617-918-1505; e-mail <weiss.jeri@epa.gov>.

Luke Trip, Environment Canada, 351 St. Joseph Blvd., Hull, Quebec K1A0H3, Canada; tel. 819-997-1967; fax 819-994-3479; e-mail <luke.trip@ec.gc.ca>.

The views expressed in this manuscript are the professional opinions of the authors and should not be interpreted as policies of their respective agencies.

**References**

1. Environmental Protection Agency (US). Executive summary. Vol. I. In: Environmental Protection Agency (US). Mercury study report to Congress. Washington: EPA; 1997. Pub. No.: EPA-452/R-97-003.
2. Agency for Toxic Substances and Disease Registry (US). Toxicological profile for mercury. Atlanta: Centers for Disease Control and Prevention; 1999.
3. Gebel T, Dunkelberg H. [Influence of chewing gum consumption and dental contact of amalgam fillings to different metal restorations on urine mercury content.] *Zentralbl Hyg Umweltmed* 1996;199:69-75.
4. Isacson G, Barregard L, Selden A, Bodin L. Impact of nocturnal bruxism on mercury uptake from dental amalgams. *Eur J Oral Sci* 1997;105:251-7.
5. Eggleston DW, Nylander M. Correlation of dental amalgam with mercury in brain tissue. *J Prosthet Dent* 1987;58:704-7.
6. Nylander M, Friberg L, Lind B. Mercury concentrations in the human brain and kidneys in relation to exposure from dental amalgam fillings. *Swed Dent J* 1987;11(5):179-87.
7. Weiner JA, Nylander M. The relationship between mercury concentration in human organs and different predictor variables. *Sci Total Environ* 1993;138(1-3):101-15.
8. Drasch G, Gath HJ, Heissler E, Schupp J, Roider G. Silver concentrations in human tissues, their dependence on dental amalgam and other factors. *J Trace Elem Med Biol* 1995;9:82-7.
9. Drexler H, Schaller KH. The mercury concentration in breast milk resulting from amalgam fillings and dietary habits. *Environ Res* 1998;77:124-9.
10. Drasch G, Aigner S, Roider G, Stalger F, Lipowsky G. Mercury in human colostrum and early breast milk: its dependence on dental amalgam and other factors. *J Trace Elem Med Biol* 1998;12:23-7.
11. Oskarsson A, Schuitz A, Skerfving S, Hallen IP, Ohlin B, Lagerkvist BJ. Total and inorganic mercury in breast milk in relation to fish consumption and amalgam in lactating women. *Arch Environ Health* 1996;51:234-41.
12. Echeverria D, Aposhian HV, Woods JS, Hayer NJ, Aposhian MM, Bittner AC Jr, et al. Neurobehavioral effects from exposure to dental amalgam Hg(0): new distinctions between recent exposure and Hg body burden. *FASEB J* 1998;12:971-80.
13. Jones DW. Exposure or absorption and the crucial question of limits for mercury. *J Can Dent Assoc* 1999;65:42-6.
14. Eley BM. The future of dental amalgam: a review of the literature. Part 5: mercury in the urine, blood and body organs from amalgam fillings. *Br Dent J* 1997;182:413-17.
15. National Institute for Occupational Safety and Health (US). A recommended standard for occupational exposure to inorganic mercury. Cincinnati (OH): NIOSH; 1977.
16. Occupational Safety and Health Administration (US). Mercury. Job Health Hazard Series. Washington: OSHA; 1975. OSHA Report No. 2234. ■



KARL BADEN